arXiv:1111.4066v1 [math.NT] 17 Nov 2011

Determinantal and Permanental Representation of Generalized Fibonacci Polynomials

Adem Sahin[a], Kenan Kaygisiz[a,*]

[a] *Department of Mathematics, Faculty of Arts and Sciences, Gaziosmanpaşa University,*
60250 Tokat, Turkey

Abstract

In this paper, we give some determinantal and permanental representations of Generalized Fibonacci Polynomials by using various Hessenberg matrices. These results are general form of determinantal and permanental representations of k sequences of the generalized order-k Fibonacci and Pell numbers.

Keywords: k sequences of the generalized order-k Fibonacci numbers, k sequences of the generalized order-k Pell numbers, generalized Fibonacci Polynomials, Hessenberg Matrix.

---

1. Introduction

Fibonacci numbers, Pell numbers and their generalizations have been studying for a long time. One of these generalizations was given by Miles in 1960.

Miles [8] defined generalized order-k Fibonacci numbers(GOkF) as,

$$f_{k,n} = \sum_{j=1}^{k} f_{k,n-j} \tag{1}$$

for $n > k \geq 2$, with boundary conditions: $f_{k,1} = f_{k,2} = f_{k,3} = \cdots = f_{k,k-2} = 0$ and $f_{k,k-1} = f_{k,k} = 1$.

Er [2] defined k sequences of generalized order-k Fibonacci numbers ($k$SO$k$F) as; for $n > 0$, $1 \leq i \leq k$

$$f_{k,n}^{i} = \sum_{j=1}^{k} c_j f_{k,n-j}^{i} \tag{2}$$

with boundary conditions for $1 - k \leq n \leq 0$,

---

[*] Corresponding author (+903562521616-3087)

Email addresses: `adem.sahin@gop.edu.tr` (Adem Sahin), `kenan.kaygisiz@gop.edu.tr` (Adem Sahin)

$$f_{k,n}^{\,i}=\begin{cases}1 & \text{if } \ i=1-n,\\ 0 & \text{otherwise,}\end{cases}$$

where c_j $(1\leq j\leq k)$ are constant coefficients, $f_{k,n}^{\,i}$ is the n-th term of i-th sequence of order-k generalization. For $c_j=1$, k-th sequence of this generalization involves the Miles generalization(1) for $i=k$, i.e.

$$f_{k,n}^{k}=f_{k,k+n-2}.$$

Kiliç and Taçcı[4] defined k sequences of generalized order-k Pell numbers ($k$SO$k$P) as; for $n>0$, $1\leq i\leq k$

$$p_{k,n}^{\,i}=2p_{k,n-1}^{\,i}+p_{k,n-2}^{\,i}+\cdots+\ p_{k,n-k}^{\,i} \tag{3}$$

with initial conditions for $1-k\leq n\leq 0$,

$$p_{k,n}^{\,i}=\begin{cases}1 & \text{if } \ i=1-n,\\ 0 & \text{otherwise,}\end{cases}$$

where $p_{k,n}^{\,i}$ is the n-th term of i-th sequence of order k generalization.

Kaygisiz and Şahin [3] defined k sequences of the generalized order-k Van der Laan numbers(kSOkV) as; for $n>0$, $1\leq i\leq k$

$$v_{k,n}^{i}=\sum_{j=2}^{k}v_{k,n-j}^{i} \tag{4}$$

with initial conditions for $1-k\leq n\leq 0$,

$$v_{k,n}^{i}=\begin{cases}1 & \text{if } i-n=k,\\ 0 & \text{otherwise}\end{cases}$$

for $1-k\leq n\leq 0$, where $v_{k,n}^{i}$ is the n-th term of i-th sequence.

MacHenry [5] defined generalized Fibonacci polynomials $(F_{k,n}(t))$ where t_i $(1\leq i\leq k)$ are constant coefficients of the core polynomial

$$P(x;t_1,t_2,\ldots,t_k)=x^k-t_1x^{k-1}-\cdots-t_k,$$

which is denoted by the vector

$$t=(t_1,t_2,\ldots,t_k).$$

$F_{k,n}(t)$ is defined inductively by

$$\begin{aligned} F_{k,n}(t) &= 0,\ n<1 \\ F_{k,1}(t) &= 1 \\ F_{k,2}(t) &= t_1 \\ F_{k,n+1}(t) &= t_1F_{k,n}(t)+\cdots+t_kF_{k,n-k+1}(t). \end{aligned} \tag{5}$$

For example the generalized Fibonacci polynomials for $k = 4$ and $k = 5$ are;

$$1, t_1, t_2 + t_1^2, t_3 + 2t_1t_2 + t_1^3, t_4 + 2t_1t_3 + t_2^2 + t_1^4 + 3t_1^2t_2, ...$$

and

$$\begin{aligned}&1,\ t_1,\ t_2 + t_1^2,\ t_3 + 2t_1t_2 + t_1^3,\ t_4 + 2t_1t_3 + t_2^2 + t_1^4 + 3t_1^2t_2,\\ &t_1^5 + 4t_1^3t_2 + 3t_1^2t_3 + 3t_1t_2^2 + 2t_1t_4 + 2t_2t_3 + t_5,\\ &2t_1t_5 + 2t_2t_4 + 6t_1t_2t_3 + t_2^3 + t_3^2 + t_1^6 + 3t_1^2t_4 + 4t_1^3t_3 + 5t_1^4t_2 + 6t_1^2t_2^2, ...\end{aligned}$$

respectively.

MacHenry studied on these polinomials and obtain very useful properties of these polynomials in [6,7].

Remark 1.1. $f_{k,n}$, $f_{k,n}^i$, $p_{k,n}^i$, $v_{k,n}^i$ and $F_{k,n}(t)$ are GOkF (1), $k$SO$k$F (2), $k$SO$k$P (3), $k$SO$k$V (4) and generalized Fibonacci polynomials (5) respectively, then

i) substituting $c_j = t_i$ in (2) and generalized Fibonacci polynomials, for $1 \leq i, j \leq k$ we obtain

$$F_{k,n}(t) = f_{k,n-1}^1,$$

ii) substituting $t_1 = 2$ and $t_i = 1$ for $2 \leq i \leq k$ in generalized Fibonacci polynomials, we obtain

$$F_{k,n}(t) = p_{k,n}^k,$$

iii) substituting $t_1 = 0$ and $t_i = 1$ for $2 \leq i \leq k$ in generalized Fibonacci polynomials, we obtain

$$F_{k,n}(t) = v_{k,n}^k,$$

iv) substituting $t_i = 1$ in generalized Fibonacci polynomials, we obtain

$$F_{k,n}(t) = f_{k,k+n-2}.$$

Minc [9] found Hessenberg matrix whose permanent are subscripted generalized order-k Fibonacci numbers. Ocal [10] gave various Hessenberg matrices whose determinants and permanents are subscripted generalized order-k Fibonacci numbers.

In this paper we derive determinantal and permanental representation of generalized Fibonacci polynomials using various Hessenberg matrices.

2. The determinantal representations

An $n \times n$ matrix $A_n = (a_{ij})$ is called lower Hessenberg matrix if $a_{ij} = 0$ when $j - i > 1$ i.e.,

$$A_n = \begin{bmatrix} a_{11} & a_{12} & 0 & \cdots & 0 \\ a_{21} & a_{22} & a_{23} & \cdots & 0 \\ a_{31} & a_{32} & a_{33} & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdots & a_{n-1,n} \\ a_{n,1} & a_{n,2} & a_{n,3} & \cdots & a_{n,n} \end{bmatrix}$$

Theorem 2.1. [1] *A_n be the $n \times n$ lower Hessenberg matrix for all $n \geq 1$ and define* $\det(A_0) = 1$, *then,*

$$\det(A_1) = a_{11}$$

and for $n \geq 2$

$$\det(A_n) = a_{n,n} \det(A_{n-1}) + \sum_{r=1}^{n-1} ((-1)^{n-r} a_{n,r} \prod_{j=r}^{n-1} a_{j,j+1} \det(A_{r-1})). \tag{6}$$

Theorem 2.2. *Let $k \geq 2$ be an integer, $F_{k,n}(t)$ be the generalized Fibonacci Polynomial (5) and $Q_{k,n} = (q_{rs})$ $n \times n$ Hessenberg matrix, where*

$$q_{rs} = \begin{cases} i^{|r-s|} . \frac{t_{r-s+1}}{t_2^{(r-s)}} & \text{if } \; -1 \leq r - s < k \;, \\ 0 & \text{otherwise} \end{cases}$$

i.e.,

$$Q_{k,n} = \begin{bmatrix} t_1 & it_2 & 0 & 0 & \cdots & 0 \\ i & t_1 & it_2 & 0 & \cdots & 0 \\ i^2 \frac{t_3}{t_2^2} & i & t_1 & it_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & & \vdots \\ i^{k-1} \frac{t_k}{t_2^{k-1}} & i^{k-2} \frac{t_{k-1}}{t_2^{k-2}} & i^{k-3} \frac{t_{k-2}}{t_2^{k-3}} & i^{k-4} \frac{t_{k-3}}{t_2^{k-4}} & \cdots & 0 \\ 0 & i^{k-1} \frac{t_k}{t_2^{k-1}} & i^{k-2} \frac{t_{k-1}}{t_2^{k-2}} & i^{k-3} \frac{t_{k-2}}{t_2^{k-3}} & \cdots & 0 \\ & \vdots & \vdots & \vdots & \ddots & \\ 0 & 0 & 0 & \cdots & \cdots & t_1 \end{bmatrix} \tag{7}$$

then

$$\det(Q_{k,n}) = F_{k,n+1}(t)$$

where $t_0 = 1$ and $i = \sqrt{-1}$.

Proof. To prove $\det(Q_{k,m}) = F_{k,m+1}(t)$, we use the mathematical induction on m. The result is true for $m = 1$ by hypothesis.

Using Theorem (2.1) we have

$$\begin{aligned}
\det(Q_{k,m+1}) &= q_{m+1,m+1}\det(Q_{k,m}) + \sum_{r=1}^{m}\left((-1)^{m+1-r}q_{m+1,r}\prod_{j=r}^{m}q_{j,j+1}\det(Q_{k,r-1})\right)\\
&= t_1\det(Q_{k,m}) + \sum_{r=1}^{m-k+1}\left((-1)^{m+1-r}q_{m+1,r}\prod_{j=r}^{m}q_{j,j+1}\det(Q_{k,r-1})\right)\\
&\quad + \sum_{r=m-k+2}^{m}\left((-1)^{m+1-r}q_{m+1,r}\prod_{j=r}^{m}q_{j,j+1}\det(Q_{k,r-1})\right)\\
&= t_1\det(Q_{k,m}) + \sum_{r=m-k+2}^{m}\left((-1)^{m+1-r}q_{m+1,r}\prod_{j=r}^{m}q_{j,j+1}\det(Q_{k,r-1})\right)\\
&= t_1\det(Q_{k,m}) + \sum_{r=m-k+2}^{m}\left((-1)^{m+1-r}.i^{m+1-r}\frac{t_{m-r+2}}{t_2^{(m-r+1)}}\prod_{j=r}^{m}it_2\det(Q_{k,r-1})\right)\\
&= t_1\det(Q_{k,m})\\
&\quad + \sum_{r=m-k+2}^{m}\left((-1)^{m+1-r}.i^{m+1-r}\frac{t_{m-r+2}}{t_2^{(m-r+1)}}.i^{m+1-r}.t_2^{(m-r+1)}\det(Q_{k,r-1})\right)\\
&= t_1\det(Q_{k,m}) + \sum_{r=m-k+2}^{m}\left((-1)^{m+1-r}.i^{m+1-r}t_{m-r+2}.i^{m+1-r}.\det(Q_{k,r-1})\right)\\
&= t_1\det(Q_{k,m}) + \sum_{r=m-k+2}^{m}t_{m-r+2}\det(Q_{k,r-1})\\
&= t_1\det(Q_{k,m}) + t_2\det(Q_{k,m-1}) + \cdots + t_k\det(Q_{k,m-(k-1)})
\end{aligned}$$

From the hypothesis and the definition of generalized Fibonacci polynomials we obtain

$$\det(Q_{k,m+1}) = t_1F_{k,m+1} + t_2F_{k,m} + \cdots + t_kF_{k,m-(k-2)} = F_{k,m+2}.$$

Therefore, the result is true for all positive integers. □

Example 2.3. *We obtain* 5*-th generalized Fibonacci polynomial for* $k = 6$*, using Theorem 2.2*

$$F_{6,5}(t) = \det\begin{bmatrix} t_1 & it_2 & 0 & 0\\ i & t_1 & it_2 & 0\\ \frac{-t_3}{t_2^2} & i & t_1 & it_2\\ \frac{-it_4}{t_2^3} & \frac{-t_3}{t_2^2} & i & t_1 \end{bmatrix} = t_4 + 2t_1t_3 + t_2^2 + t_1^4 + 3t_1^2t_2.$$

Corollary 2.4. [10] *Let* $k \geq 2$ *be an integer,* $f_{k,n}$ *be the generalized order-k Fibonacci numbers (1) and* $C_{k,n} = (c_{rs})$ $n \times n$ *Hessenberg matrix, where*

$$c_{rs} = \begin{cases} i^{|r-s|} & \textit{if} \quad -1 \leq r-s < k \ , \\ 0 & \textit{otherwise} \end{cases}$$

then

$$\det(C_{k,n}) = f_{k,k+n-1}$$

where $i = \sqrt{-1}$.

Proof. It is direct from Theorem 2.2 for $t_i = 1$. □

Theorem 2.5. *Let* $k \geq 2$ *be an integer,* $F_{k,n}(t)$ *be the generalized Fionacci Polynomial (5) and* $B_{k,n} = (b_{ij})$ *be an* $n \times n$ *lower Hessenberg matrix such that*

$$b_{ij} = \begin{cases} -t_2 & \textit{if} \quad j = i+1, \\ \frac{t_{i-j+1}}{t_2^{(i-j)}} & \textit{if} \quad 0 \leq i-j < k, \\ 0 & \textit{otherwise} \end{cases}$$

i.e.,

$$B_{k,n} = \begin{bmatrix} t_1 & -t_2 & 0 & 0 & \cdots & 0 \\ 1 & t_1 & -t_2 & 0 & \cdots & 0 \\ \frac{t_3}{t_2^2} & 1 & t_1 & -t_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & & \vdots \\ \frac{t_k}{t_2^{k-1}} & \frac{t_{k-1}}{t_2^{k-2}} & \frac{t_{k-2}}{t_2^{k-3}} & \frac{t_{k-3}}{t_2^{k-4}} & \cdots & 0 \\ 0 & \frac{t_k}{t_2^{k-1}} & \frac{t_{k-1}}{t_2^{k-2}} & \frac{t_{k-2}}{t_2^{k-3}} & \cdots & 0 \\ & \vdots & \vdots & \vdots & \ddots & \\ 0 & 0 & 0 & \cdots & \cdots & t_1 \end{bmatrix} \tag{8}$$

then

$$\det(B_{k,n}) = F_{k,n+1}(t).$$

where $t_0 = 1$.

Proof. To prove $\det(B_{k,m}) = F_{k,m+1}(t)$, we use the mathematical induction on m. The result is true for $m = 1$ by hypothesis.

Using Theorem (2.1) we have

$$\begin{aligned}
\det(B_{m+1,k}) &= b_{m+1,m+1}\det(B_{k,m}) + \sum_{r=1}^{m}((-1)^{m+1-r}b_{m+1,r}\prod_{j=r}^{m} b_{j,j+1}\det(B_{r-1,k})) \\
&= t_1\det(B_{k,m}) + \sum_{r=1}^{m-k+1}((-1)^{m+1-r}b_{m+1,r}\prod_{j=r}^{m} b_{j,j+1}\det(B_{r-1,k})) \\
&\quad + \sum_{r=m-k+2}^{m}((-1)^{m+1-r}b_{m+1,r}\prod_{j=r}^{m} b_{j,j+1}\det(B_{r-1,k})) \\
&= t_1\det(B_{k,m}) + \sum_{r=m-k+2}^{m}((-1)^{m+1-r}.\frac{t_{m-r+2}}{t_2^{(m-r+1)}}\prod_{j=r}^{m}(-t_2)\det(B_{r-1,k})) \\
&= t_1\det(B_{k,m}) \\
&\quad + \sum_{r=m-k+2}^{m}((-1)^{m+1-r}.\frac{t_{m-r+2}}{t_2^{(m-r+1)}}.(-1)^{m+1-r}t_2^{(m-r+1)}\det(B_{r-1,k})) \\
&= t_1\det(B_{k,m}) + \sum_{r=m-k+2}^{m}(t_{m-r+2}.\det(B_{r-1,k})) \\
&= t_1\det(B_{k,m}) + t_2\det(B_{k,m-1}) + \cdots + t_k\det(B_{k,m-(k-1)}).
\end{aligned}$$

From the hypothesis and the definition of generalized Fibonacci polynomials we obtain

$$\det(Q_{m+1,k}) = t_1F_{k,m+1} + t_2F_{k,m} + \cdots + t_kF_{k,m-(k-2)} = F_{k,m+2}.$$

Therefore, the result is true for all positive integers. $\square$

Example 2.6. *We obtain* 6*-th generalized Fibonacci polynomial for* $k = 4$*, using Theorem 2.5*

$$F_{4,6}(t) = \det\begin{bmatrix} t_1 & -t_2 & 0 & 0 & 0 \\ 1 & t_1 & -t_2 & 0 & 0 \\ \frac{t_3}{t_2^2} & 1 & t_1 & -t_2 & 0 \\ \frac{t_4}{t_2^3} & \frac{t_3}{t_2^2} & 1 & t_1 & -t_2 \\ 0 & \frac{t_4}{t_2^3} & \frac{t_3}{t_2^2} & 1 & t_1 \end{bmatrix} = 2t_1t_4 + 2t_2t_3 + t_1^5 + 3t_1t_2^2 + 3t_1^2t_3 + 4t_1^3t_2.$$

Corollary 2.7. [10]*Let* $k \geq 2$ *be an integer,* $f_{k,n}$ *be the generalized order-k Fibonacci numbers (1) and* $M_{k,n} = (m_{ij})$ *be an* $n \times n$ *lower Hessenberg matrix such that*

$$m_{ij} = \begin{cases} -1 & \text{if} \quad j = i+1, \\ 1 & \text{if} \quad 0 \leq i-j < k, \\ 0 & \text{otherwise} \end{cases}$$

then

$$\det(M_{k,n}) = f_{k,k+n-1}$$

Proof. It is direct from Theorem 2.5 for $t_i = 1$. $\square$

Corollary 2.8. *If we rewrite Theorem (2.2) and Theorem (2.5)*
i) for $t_i = c_j$ $(1 \leq i, j \leq k)$, *we obtain*

$$\det(Q_{k,n}) = f^1_{k,n}$$

and

$$\det(B_{k,n}) = f^1_{k,n}$$

respectively,
ii) for $t_1 = 2$ *and* $t_i = 1$ *for* $2 \leq i \leq k$, *we obtain*

$$\det(Q_{k,n}) = p^k_{k,n+1}$$

and

$$\det(B_{k,n}) = p^k_{k,n+1}$$

respectively,
iii) for $t_1 = 0$ *and* $t_i = 1$ *for* $2 \leq i \leq k$, *we obtain*

$$\det(Q_{k,n}) = v^k_{k,n+1}$$

and

$$\det(B_{k,n}) = v^k_{k,n+1}$$

respectively. Where $f^1_{k,n}$, $p^k_{k,n}$, $v^k_{k,n}$ *be the k sequences of generalized order-k Fibonacci, Pell and Van der Laan numbers. Matrices* $Q_{k,n}$ *and* $B_{k,n}$ *are as in (7) and (8), respectively.*

3. The permanent representations

Let $A = (a_{i,j})$ be a square matrix of order n over a ring R. The permanent of A is defined by

$$\operatorname{per}(A) = \sum_{\sigma \in S_n} \prod_{i=1}^{n} a_{i,\sigma(i)}$$

where S_n denotes the symmetric group on n letters.

Let $A_{i,j}$ be the (i,j)-th minor of matrix A. Then

$$\operatorname{per}(A) = \sum_{k=1}^{n} a_{i,k} \operatorname{per}(A_{i,k}) = \sum_{k=1}^{n} a_{k,j} \operatorname{per}(A_{k,j})$$

for any i, j.

Theorem 3.1. [10]*Let A_n be $n \times n$ lower Hessenberg matrix for all $n \geq 1$ and define $per(A_0) = 1$. Then,*

$$per(A_1) = a_{11}$$

and for $n \geq 2$

$$per(A_n) = a_{n,n} per(A_{n-1}) + \sum_{r=1}^{n-1} (a_{n,r} \prod_{j=r}^{n-1} a_{j,j+1} per(A_{r-1})). \tag{9}$$

Theorem 3.2. *Let $k \geq 2$ be an integer, $F_{k,n}(t)$ be the generalized Fibonacci Polynomial and $H_{k,n} = (h_{rs})$ be an $n \times n$ lower Hessenberg matrix such that*

$$h_{rs} = \begin{cases} i^{r-s} . \frac{t_{r-s+1}}{t_2^{(r-s)}} & \text{if } \ -1 \leq r-s < k \\ 0 & \text{otherwise} \end{cases}$$

i.e.,

$$H_{k,n} = \begin{bmatrix} t_1 & -it_2 & 0 & 0 & \cdots & 0 \\ i & t_1 & -it_2 & 0 & \cdots & 0 \\ i^2 \frac{t_3}{t_2^2} & i & t_1 & -it_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & & \vdots \\ i^{k-1} \frac{t_k}{t_2^{k-1}} & i^{k-2} \frac{t_{k-1}}{t_2^{k-2}} & i^{k-3} \frac{t_{k-2}}{t_2^{k-3}} & i^{k-4} \frac{t_{k-3}}{t_2^{k-4}} & \cdots & 0 \\ 0 & i^{k-1} \frac{t_k}{t_2^{k-1}} & i^{k-2} \frac{t_{k-1}}{t_2^{k-2}} & i^{k-3} \frac{t_{k-2}}{t_2^{k-3}} & \cdots & 0 \\ & \vdots & \vdots & \vdots & \ddots & \\ 0 & 0 & 0 & \cdots & \cdots & t_1 \end{bmatrix} \tag{10}$$

then

$$per(H_{k,n}) = F_{k,n+1}(t)$$

where $t_0 = 1$ and $i = \sqrt{-1}$.

Proof. Since the proof is similar to the proof of Theorem (2.2) by using Theorem (3.1) we omit the detail. □

Example 3.3. *We obtain 7-th generalized Fibonacci Polynomials for $k = 5$, using Theorem 3.2*

$$\begin{aligned} F_{5,7} &= per \begin{bmatrix} t_1 & -it_2 & 0 & 0 & 0 & 0 \\ i & t_1 & -it_2 & 0 & 0 & 0 \\ \frac{-t_3}{t_2^2} & i & t_1 & -it_2 & 0 & 0 \\ \frac{-it_4}{t_2^3} & \frac{-t_3}{t_2^2} & i & t_1 & -it_2 & 0 \\ \frac{t_5}{t_2^4} & \frac{-it_4}{t_2^3} & \frac{-t_3}{t_2^2} & i & t_1 & -it_2 \\ 0 & \frac{t_5}{t_2^4} & \frac{-it_4}{t_2^3} & \frac{-t_3}{t_2^2} & i & t_1 \end{bmatrix} \\ &= 2t_1t_5 + 2t_2t_4 + 6t_1t_2t_3 + t_2^3 + t_3^2 + t_1^6 + 3t_1^2t_4 + 4t_1^3t_3 + 5t_1^4t_2 + 6t_1^2t_2^2. \end{aligned}$$

Corollary 3.4. [10]*Let $k \geq 2$ be an integer, $f_{k,n}$ be the generalized order-k Fibonacci numbers and $H_{k,n} = (h_{rs})$ be an $n \times n$ lower Hessenberg matrix such that*

$$h_{rs} = \begin{cases} i^{r-s} & \text{if } \; -1 \leq r-s < k \\ 0 & \text{otherwise} \end{cases}$$

then

$$per(H_{k,n}) = f_{k,k+n-1}$$

Proof. It is direct from Theorem 3.2 for $t_i = 1$. □

Theorem 3.5. *Let $k \geq 2$ be an integer, $F_{k,n}(t)$ be the generalized Fibonacci Polynomial and $L_{k,n} = (l_{ij})$ be an $n \times n$ lower Hessenberg matrix such that*

$$l_{ij} = \begin{cases} \frac{t_{i-j+1}}{t_2^{(i-j)}} & \text{if } \; -1 \leq i-j < k, \\ 0 & \text{otherwise} \end{cases}$$

i.e.,

$$L_{k,n} = \begin{bmatrix} t_1 & t_2 & 0 & 0 & \cdots & 0 \\ 1 & t_1 & t_2 & 0 & \cdots & 0 \\ \frac{t_3}{t_2^2} & 1 & t_1 & t_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & & \vdots \\ \frac{t_k}{t_2^{k-1}} & \frac{t_{k-1}}{t_2^{k-2}} & \frac{t_{k-2}}{t_2^{k-3}} & \frac{t_{k-3}}{t_2^{k-4}} & \cdots & 0 \\ 0 & \frac{t_k}{t_2^{k-1}} & \frac{t_{k-1}}{t_2^{k-2}} & \frac{t_{k-2}}{t_2^{k-3}} & \cdots & 0 \\ & \vdots & \vdots & \vdots & \ddots & \\ 0 & 0 & 0 & \cdots & \cdots & t_1 \end{bmatrix}. \tag{11}$$

where $t_0 = 1$, then

$$per(L_{k,n}) = F_{k,n+1}.$$

Proof. This is similar to the proof of Theorem 2.5 using Theorem 3.1. □

Corollary 3.6. [9]*Let $k \geq 2$ be an integer, $f_{k,n}$ be the generalized order-k Fibonacci numbers and $D_{k,n} = (d_{ij})$ be an $n \times n$ lower Hessenberg matrix such that*

$$d_{ij} = \begin{cases} 1 & \text{if} \quad -1 \leq i-j < k, \\ 0 & \text{otherwise} \end{cases}$$

then

$$per(D_{k,n}) = f_{k,k+n-1}.$$

Proof. It is direct from Theorem 3.5 for $t_i = 1$. □

Corollary 3.7. *If we rewrite Theorem (3.2) and Theorem (3.5)*
i) for $t_i = c_j$ $(1 \leq i, j \leq k)$, we obtain

$$per(H_{k,n}) = f_{k,n}^{1}$$

and

$$per(L_{k,n}) = f_{k,n}^{1}$$

respectively,
ii) for $t_1 = 2$ and $t_i = 1$ $(2 \leq i \leq k)$, we obtain

$$per(H_{k,n}) = p_{k,n+1}^{k}$$

and

$$per(L_{k,n}) = p_{k,n+1}^{k}$$

respectively,
iii) for $t_1 = 0$ and $t_i = 1$ $(2 \leq i \leq k)$, we obtain

$$per(H_{k,n}) = v_{k,n+1}^{k}$$

and

$$per(L_{k,n}) = v_{k,n+1}^{k}$$

respectively. Where $f_{k,n}^{k}$, $p_{k,n}^{k}$, $v_{k,n}^{k}$ be the k sequences of generalized order-k Fibonacci, Pell and Van der Laan numbers. Matrices $H_{k,n}$ and $L_{k,n}$ are as in (10) and (11), respectively.

References

[1] N.D. Cahill, J.R. D'Errico, D.A. Narayan, J.Y. Narayan, Fibonacci determinants, College Math. J. 33(3) (2002) 221-225.

[2] M. C. Er, Sums of Fibonacci Numbers by Matrix Method, Fibonacci Quarterly, vol.23, no. 3, 204-207, (1984).

[3] K. Kaygisiz and A. Şahin, k sequences of Generalized Van der Laan and Generalized Perrin Polynomials. *to submitted*.

[4] E. Kılıç and D. Taşci, The Generalized Binet Formula, Representation and Sums of The Generalized Order-k Pell Numbers. *Taiwanese Jour. of Math.* 10(2006), no. 6, 1661-1670.

[5] T. MacHenry, A Subgroup of The Group of Units in The Ring of Arithmetic Fonctions. *Rocky Mount. Jour. of Math.*39(1999), no. 3, 1055-1065.

[6] T. MacHenry, Generalized Fibonacci and Lucas Polynomials and Multiplicative Arithmetic Functions. *Fibonacci Quarterly.* 38(2000), 17-24.

[7] T. MacHenry and K. Wong, Degree k Linear Recursions mod(p) and Number Fields. *Rocky Mount. Jour. of Math. to appear.*

[8] E.P. Miles, Generalized Fibonacci Numbers and Associated Matrices. Amer. Math. Monthly 67 (1960), 745-752.

[9] H. Minc, Encyclopaedia of Mathematics and its Applications, Permanents, Vol.6, Addison-Wesley Publishing Company, London, 1978.

[10] A.A. Öcal, N. Tuglu, E. Altinisik, On the representation of k-generalized Fibonacci and Lucas Numbers. Appl. Math. and Comput. 170(2005) 584-596.